

Pronto Plus Kill Bedbugs and Dust Mites

ACTIVE INGREDIENTS:

3-phenoxybenzyl-(1RS,3RS, 1RS, 3SR)-2, 2-dimethyl-3-(2-methylprop-1-enyl)

cyclopropanecarboxylate 0.4%

N-Octyl bicycloheptene dicarboximide. 1.6%

OTHER INGREDIENTS: 98.0%
100.0%

STOP: READ ENTIRE LABEL BEFORE USE

KEEP OUT OF REACH OF CHILDREN

CAUTION

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. **USE RESTRICTIONS:** In hospitals and nursing homes, remove patients before treatment. Ventilate room for two hours before returning patients. This product is not for use on humans: Do not use in commercial food/feed processing, preparation, food/feed storage or serving areas. In the home, all food/feed processing surfaces and utensils should be covered during treatment, or thoroughly washed before use. Exposed food/feed should be covered or removed. Remove pets, birds and cover fish aquariums before spraying.

SHAKE WELL BEFORE EACH USE. APPLICATION: Spray in an inconspicuous area to test for possible staining or discoloration. Evaluate test area after drying. Hold container upright with nozzle aimed away from you. Depress valve and spray from a distance of 8 to 10 inches. Spray each square foot for 5 seconds or until damp. Most effective results are achieved when used as part of a treatment protocol that includes physical, environmental and other chemical pest control measures. Vacate room after treatment and ventilate before reoccupying. Do not allow children or pets to contact treated areas until surfaces are dry. Allow sprayed articles and surfaces to dry thoroughly before using or replacing bedding.

TO KILL DUST MITES: Thoroughly vacuum entire room concentrating on areas where mites congregate: such as mattresses, box springs, headboards, walls, floors, carpeting, and baseboards. Then spray surfaces until damp.

TO KILL BEDBUGS and BEDBUG EGGS: Apply as a spot treatment to cracks and crevices on and around baseboards, floorboards, bed frames, wall hangings, headboards, furniture, door and window frames, millwork and walls. Apply as a surface spray to carpet, mattresses, box springs, walls, furniture, bedding, floor and floor coverings, rugs, garments, luggage, closets, drapes and other window appointments. Thoroughly clean and air-out mattresses and box springs. Then treat areas that may harbor bedbugs or their eggs such as tufts, folds, and seams. When treating for bedbug eggs, spray each square foot for 13 seconds or until damp.

STORAGE AND DISPOSAL Do not contaminate water, food, or feed by storage or disposal. **Pesticide Storage:** Store in a cool dry area. Always store pesticides in the original container. Store away from food and pet food. **Pesticide Disposal and Container Handling:** Do not puncture or incinerate! Nonrefillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION: Harmful if absorbed through skin. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse. **PHYSICAL OR CHEMICAL HAZARDS Contents under pressure.** Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting. Do not apply this product around electrical equipment due to the possibility of shock hazard.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a Poison Control Center or doctor for treatment advice.

IF SWALLOWED: Immediately call a Poison Control Center or doctor. Do not induce vomiting unless told to do so by a Poison Control Center or a doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person. Have the product container or label with you when calling a Poison Control Center or doctor, or going for treatment. For information regarding medical emergencies or pesticide incidents, call 1-888-542-3546.

EPA Reg. No. 1021-1767-75406 EPA Est. No. 5693-MA-1